

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/304032530>

nirS-encoding denitrifier community composition, distribution, and abundance along the coastal wetlands of China

Article in *Applied Microbiology and Biotechnology* · June 2016

DOI: 10.1007/s00253-016-7659-5

CITATIONS

3

READS

134

11 authors, including:



Juan Gao

East China Normal University

17 PUBLICATIONS 71 CITATIONS

SEE PROFILE



Lijun Hou

East China Normal University

98 PUBLICATIONS 1,323 CITATIONS

SEE PROFILE



Yanling Zheng

East China Normal University

28 PUBLICATIONS 234 CITATIONS

SEE PROFILE



Xiaofei Li

East China Normal University

33 PUBLICATIONS 162 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



nitrogen cycling [View project](#)



PAHs research [View project](#)

nirS-Encoding denitrifier community composition, distribution, and abundance along the coastal wetlands of China

Juan Gao¹ · Lijun Hou¹ · Yanling Zheng^{1,2} · Min Liu² · Guoyu Yin^{1,2} · Xiaofei Li² · Xianbiao Lin² · Chendi Yu¹ · Rong Wang¹ · Xiaofen Jiang¹ · Xiuru Sun²

Received: 7 April 2016 / Revised: 29 May 2016 / Accepted: 1 June 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract For the past few decades, human activities have intensively increased the reactive nitrogen enrichment in China's coastal wetlands. Although denitrification is a critical pathway of nitrogen removal, the understanding of denitrifier community dynamics driving denitrification remains limited in the coastal wetlands. In this study, the diversity, abundance, and community composition of *nirS*-encoding denitrifiers were analyzed to reveal their variations in China's coastal wetlands. Diverse *nirS* sequences were obtained and more than 98 % of them shared considerable phylogenetic similarity with sequences obtained from aquatic systems (marine/estuarine/coastal sediments and hypoxia sea water). Clone library analysis revealed that the distribution and composition of *nirS*-harboring denitrifiers had a significant latitudinal differentiation, but without a seasonal shift. Canonical correspondence analysis showed that the community structure of *nirS*-encoding denitrifiers was significantly related to temperature and ammonium concentration. The *nirS* gene abundance ranged from 4.3×10^5 to 3.7×10^7 copies g^{-1} dry sediment, with a significant spatial heterogeneity. Among all detected

environmental factors, temperature was a key factor affecting not only the *nirS* gene abundance but also the community structure of *nirS*-type denitrifiers. Overall, this study significantly enhances our understanding of the structure and dynamics of denitrifying communities in the coastal wetlands of China.

Keywords Denitrification · *nirS* gene · Coastal wetlands · Temperature · Abundance · Community structure

Introduction

Over the past several decades, reactive nitrogen (Nr) input has incredibly increased because of human activities (Galloway and Cowling 2002; Galloway et al. 2008). By 2050, the nitrogen delivery in most regions is predicted to reach $50 \text{ kg N ha}^{-1} \text{ year}^{-1}$ to afford the still-growing human population (Galloway et al. 2004). The acceleration of Nr is attributed mainly to industrial production of nitrogen fertilizer and combustion of fossil fuel which contribute about $1.48 \times 10^{13} \text{ mol N year}^{-1}$ (Canfield et al. 2010; Vitousek et al. 1997). Although the increasing input of Nr has enabled humans to greatly improve crop yield, considerable Nr is eventually transported into estuarine and coastal wetlands through river flow, groundwater discharge and atmospheric deposition (Boyer et al. 2006; Seitzinger 2008), which has caused a series of environmental problems such as aquatic eutrophication, harmful algal bloom and hypoxia (Earl et al. 2006; Gruber and Galloway 2008; Howarth 2008). Therefore, it is of significance to study the nitrogen removal and associated microbial mechanisms for protecting and improving water quality in estuarine and coastal environments.

Denitrification, the dissimilatory reduction of oxidized N compounds (NO_3^- and NO_2^-) to gaseous nitrogen (including

Juan Gao, Lijun Hou, Yanling Zheng contributed equally to this work.

Electronic supplementary material The online version of this article (doi:10.1007/s00253-016-7659-5) contains supplementary material, which is available to authorized users.

✉ Lijun Hou
ljhou@sklec.ecnu.edu.cn

✉ Min Liu
mliu@geo.ecnu.edu.cn

¹ State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062, China

² College of Geographical Sciences, East China Normal University, Shanghai 200241, China

NO, N₂O, and N₂), has been considered an effective microbial nitrogen removal process (Falkowski et al. 2008; Zumft 1997). This process contributes more than 70 % of the nitrogen loss in natural ecosystems, which is much higher than anammox process (Babbin et al. 2014; Dalsgaard et al. 2012). Thus denitrification plays a crucial role in controlling nitrogen fate (van Breemen et al. 2002). Diverse types of metabolic enzymes, including nitrate reductases (Nar), nitrite reductases (Nir), nitric oxide reductases (Nor), and nitrous oxide reductases (Nos), catalyze the denitrification process (Zumft 1997). In particular, Nir catalyzes the rate-limiting step in denitrification, which is encoded by *nirK* and/or *nirS* genes (Braker et al. 2001). Although these two genes are structurally different, enzyme types are functionally and physiologically similar (Coyne et al. 1989). Compared with *nirK* which is also contained in nitrifiers (Cantera and Stein 2007), *nirS* is found to be more widely distributed in natural environments (Braker et al. 1998). Therefore, *nirS* has been most frequently used for functional biomarkers of denitrifying community.

China has contributed far more Nr than other countries, due to large population and rapid agricultural and economic development (Cui et al. 2013). Over the past century, Nr emission had increased from 9.2 to 56 Tg year⁻¹ in China (Cui et al. 2013), and the majority of the Nr is delivered into the environment (Galloway and Cowling 2002; Wang and Wang 2009). At present, the estuaries and coastal seas of China have greatly suffered from the severe Nr pollution (Cui et al. 2013; Zhao et al. 2012). Although the important role of denitrification in the nitrogen removal is identified, little is known about the dynamics of denitrifiers in the nitrogen-enriched environments. The objectives of this study are to investigate the abundance, diversity, and distribution of *nirS*-harboring denitrifiers along the coastal wetlands of China and to explore the underlying interactions among the dynamics of denitrifiers, environmental parameters, and denitrifying activities in the wetland ecosystem.

Materials and methods

Study area

The coastal regions of China cover an area over three million square kilometers, with a 32,000 km coastline (Wang 1992). In China, the development of coastal regions has made a significant contribution to the national economy, harboring more than 60 % of China's gross domestic product (He et al. 2014). Over the past two decades, the coastal wetland areas have decreased 3.38×10^6 ha, with a total loss rate of 9.33 % (Wetland China 2014). The loss of the coastal wetlands is mainly attributed to the increasing growth of population in the coastal zone, which was coupled with rising coastal urban areas, economic growth, rapid urbanization, and infrastructure development (He et al. 2014; Shi et al. 2015). In addition, the

interference of human activities has caused a series of ecological changes in China's coastal wetland ecosystems, such as the losses of biological habitat, productivity, and diversity (Lin et al. 2007; Sun et al. 2015; Xie et al. 2010).

Sample collection

In this study, sediment samples were collected from eleven sites (P1 to P11) located in the coastal wetlands of China (including Dandong, Tangshan, Weifang, Qingdao, Lianyungang, Shanghai, Wenzhou, Fuzhou, Shantou, Zhuhai, and Beihai) (Fig. 1), ranging from high latitude sites (P1 to P6) to low latitude sites (P7 to P11). The numbering scheme for sampling sites decreases with latitude. Field work was conducted in winter (January) and summer (August) 2014, respectively. Triplicate sediment samples (0–5 cm) were collected with PVC cores at each site. The core sediments were transported to the lab on ice and immediately homogenized as one composite sample under helium condition. A fraction of the composite sample was archived at -80 °C until DNA isolation and used for microbial molecular analysis. The remaining sample was stored at 4 °C for analyses of sediment physicochemical parameters and denitrification activity. The data on sediment properties and denitrification rates in these collected samples have been given in Hou et al. (2015) (Table S1 and Fig. S1).

PCR amplification, cloning, and phylogenetic analysis

Total genomic DNA was extracted from ~0.3 g sediment with PowerSoil DNA isolation kits (MoBio, USA). The *nirS* gene fragment (encoding cytochrome *cd1*-containing nitrite reductase ~840–890 bp) was amplified from sediment DNA extracts using primers *nirS*-1F and *nirS*-6R (Braker et al.




Fig. 1 Location of sampling sites within the coastal wetlands of China

1998). The details of the primers and PCR conditions are shown in Table S2. The amplification products were visualized by electrophoresis on 1.0 % agarose gels. The obtained PCR products were purified using the Gel Advance gel extraction system (Viogene, China), and cloned into the pUCm-T Vector (Sangon, China). Then insert-containing transformants were transformed into *E. coli* XL1-Blue for growth. Approximate 100 clones were randomly screened from each sample and sequenced with an ABI 3370XL Prism genetic analyzer (Applied Biosystems, Canada).

Sequences were assembled, edited and put in order with DNASTar Lasergene (DNASTar, USA). The *nirS* sequences were analyzed in GenBank by BLAST (<http://blast.st.va.ncbi.nlm.nih.gov/Blast.cgi>) to obtain reference sequences. All of the obtained sequences were aligned with the ClustalX software (Thompson et al. 1997). The Mothur program (<http://www.mothur.org/>) was used to classify the sequences into one operational taxonomic unit (OTU) with >95 % identity. Phylogenetic tree was created by Mega 5 program with the neighbor-joining method (Kumar et al. 2004). The reliability of the tree topologies was estimated by performing 1000 bootstrapping replicates (Tamura et al. 2007).

The unique nucleotide sequences obtained in this study are available, and have been deposited in GenBank database under accession numbers of KU995338 to KU996352.

qPCR of *nirS* gene

The abundance of *nirS* gene fragment (425 bp) in sediments was quantified in triplicate with primers cd3aF and R3cd (Throback et al. 2004). The details of the primers and PCR conditions are shown in Table S2. Plasmids containing cloned *nirS* PCR amplicons were generated with *Escherichia coli* hosts by using Qiagen Miniprep Spin Kit. The plasmids were diluted into a series of gradient as standard curves. The concentration of original plasmid was estimated with a Nanodrop-2000 Spectrophotometer (Thermo, USA). Quantification standard curves were formed via plating the threshold cycle (Ct) versus the log₁₀ values of *nirS* gene copy numbers carried by the standard plasmids solutions, with strong linear relationship ($R^2 = 0.9996$) and high amplification efficiency (94.7 % in winter and 100.1 % in summer). The standard curve ranged over 7 orders of magnitude of the standard plasmid's concentrations (1.71×10^3 to 1.71×10^9 copies per microliter). The melt curve for standards and samples only had a single peak at 85.4 °C, indicating that the fluorescent signals were obtained from specific DNA samples in all the process of quantitative PCR.

Statistics analysis

The biodiversity indicators (Shannon-Weiner and Simpson), species richness Chao 1 estimator, and the rarefactions curves were obtained with the Mothur program (Schloss et al. 2009).

The coverage for each constructed clone library was calculated using the follow method: the obtained OTUs number divided by chao1 indicator (Mohamed et al. 2010). Relationships between denitrifying bacterial community compositions and environmental indices were analyzed by the Canoco (version 4.5) software using canonical correspondence analysis (CCA) on the basis of the results of detrended correspondence analysis (DCA) (Danovaro and Gambi 2002; ter Braak 1988). Community classifications of sediment *nirS*-based denitrifiers were explored with principal coordinates analysis (PCoA) by the Qiime 1.9.0 software (Caporaso et al. 2010). Pearson correlation analyses were conducted with SPSS (version 16.0) software to explore correlations of environmental variables with the richness and abundance of *nirS*-harboring denitrifiers. Additionally, one-way analysis of variance (ANOVA) was performed to compare spatial and seasonal differences in *nirS*-encoding denitrifiers.

Results

nirS-based denitrifier diversity

The *nirS* gene sequences were successfully recovered from sediment DNA extracts at all sampling sites. Clone libraries were constructed for each site, containing 73 to 96 clones per library (Table 1). This resulted in a total database of 1942 *nirS* gene clones (Table 1). To date, these represented the most widespread *nirS* gene clone library in China's coastal wetlands. To analyze the diversity of *nirS*-based denitrifiers, 5 % divergence at the nucleotide level was used for *nirS* sequences to define OTU. In this study, 13 to 33 OTUs were obtained in each individual clone library (Table 1), and a total of 493 OTUs were identified. The library coverage was estimated at between 91.5–99.2 %. The high coverage indicates that the majority of the *nirS*-based denitrifier diversity was obtained, which is further confirmed by the rarefaction analysis (Fig. 2).

Based on the Shannon-Weiner and Simpson indices (Table 1), the maximal *nirS*-type denitrifier richness was found at site P6 in summer and P1 in winter where 30 and 33 OTUs were observed, respectively. The lack of significant curvature (Fig. 2) indicated that the richness of distinct *nirS*-based sequences was not yet saturated in those two libraries. The second high richness of *nirS*-type denitrifiers appeared at site P10 in summer where 28 OTUs were obtained. Additionally, relatively low biodiversity of *nirS* gene was recorded at site P8 in summer where only 13 OTUs were found. The other 18 clone libraries had intermediate diversity, with an average Shannon-Weiner index of 2.58. Overall, the diversity of *nirS* gene showed a significant spatial difference (one-way ANOVA, $P = 0.001$) along the coastal wetlands of China. However, no distinctive seasonal shift was found between

Table 1 Diversity estimators of *nirS*-encoding denitrifiers in the coastal wetlands of China

Seasons	Sites	No. of clones	OTUs ^a	Shannon ^b	Chao 1 ^c	1/Simpson ^d	Coverage (%) ^e
winter	P1	85	33	3.17	36.1	21.12	91.5
	P2	86	15	1.94	15.9	4.32	94.6
	P3	93	20	2.59	21.3	11.50	94.1
	P4	90	20	2.20	21.4	5.40	93.6
	P5	87	25	2.88	25.8	15.52	97.0
	P6	96	19	2.50	20.2	9.01	94.1
	P7	85	18	2.19	19.1	5.61	94.2
	P8	94	15	2.04	15.1	5.27	99.2
	P9	90	26	2.84	27.2	12.75	95.8
	P10	88	23	2.90	23.7	18.23	97.2
	P11	89	30	3.00	30.5	14.50	98.5
Summer	P1	88	23	2.71	23.5	11.60	98.0
	P2	94	25	2.89	25.8	16.43	96.8
	P3	87	20	2.45	20.6	7.68	97.1
	P4	90	26	2.89	26.4	14.56	98.6
	P5	73	24	2.93	25.0	16.95	96.0
	P6	79	30	3.23	31.4	29.34	95.5
	P7	89	14	1.68	14.1	2.84	99.0
	P8	93	13	1.28	13.4	1.97	97.2
	P9	91	23	2.72	24.1	11.31	95.4
	P10	85	28	3.06	28.6	19.40	97.8
	P11	90	23	2.66	24.4	11.06	94.4

^a OTUs are defined based on 5 % nucleotide acid divergence

^b Shannon–Weiner index. Higher number represents more diversity

^c Nonparametric statistical predictions of total richness of OTUs based on distribution of singletons and doubletons

^d Reciprocal of Simpson's diversity index. Higher number represents more diversity

^e Percentage of observed number of OTUs divided by Chao1 estimate

summer and winter (one-way ANOVA, $P = 0.919$), with average Shannon–Weiner indices of 2.59 and 2.57, respectively.

Community composition and distribution of *nirS*-encoding bacteria

For the phylogenetic analysis, the *nirS* gene sequences were grouped into 10 distinctly defined clusters (I to X) on the basis of evolutionary distance (Fig. 3). In this study, the levels of nucleotide clone identity ranged from 45 % to 100 %. The 1942 *nirS* gene sequences were aligned with representative database sequences and had high degree of identity (93.0–100.0 %) to the closest matched sequences retrieved from the GenBank. Phylogenetic analysis showed that a large proportion of the unique OTUs matched with uncultured environmental *nirS* assemblages. The majority of the closely matched sequences in the GenBank were retrieved from sediment environments, including Haihe River estuary (KC106788), Yangtze estuary (KF363218; EU236006;

KM892169; EU235754) (Zhang et al. 2014; Zheng et al. 2015), Bahia del Tobari estuary (KC614247), San Francisco Bay estuary (GQ453730) (Mosier and Francis 2010), Bohai Gulf (JN257854), East Pacific sea (GU348415), coastal sea in France (KJ640012) (Stauffert et al. 2014), as well as coastal regions in Baltic sea (DQ072204) (Hannig et al. 2006) and the South Pacific (AJ811468) (Castro-Gonzalez et al. 2005). The *nirS* gene sequences of cluster IX were closely related to the cultivated denitrifiers of *Proteobacteria Pseudomonadales* (CP000304) (90 % identity) (Yan et al. 2008).

All of the ten clusters corresponded to distinct groups in the phylogenetic tree (Fig. 3). The cluster II contained the maximal amount of *nirS* gene sequences, occupying 26.6 % of the total sequences (Fig. 4). However, the cluster V only accounted for 0.6 % (Fig. 4). The clusters I, II, and IV were discovered in all the 22 built clone libraries, including 1.1 % to 65.2 % of *nirS* gene sequences. Of these ten clusters, no one was unique to a particular site, and all the clusters included sequences from four or more samples. The *nirS* sequences




Fig. 2 Rarefaction analysis of *nirS*-encoding denitrifier communities in summer and winter, respectively. OTUs were defined by <5 % divergence in nucleotide sequence

from high latitude habitats (P1 to P6) were the dominant type in the clusters II, V, VIII, and X (Table S3), accounting for 61.0 % to 83.3 % of the *nirS*-based denitrifiers. However, in the clusters VI and IX, the *nirS* sequences were co-dominated by the low latitude sites (P7 to P11), accounting for 71.1 % to 84.8 %. Among all the 22 clone libraries, no one library was dispersed throughout the phylogenetic tree, which generally occupied 5 to 8 clusters.

Spatial and temporal distributions of *nirS*-harboring denitrifier community were statistically compared with the weighted UniFrac PCoA analysis (Fig. 5). The first two principal coordinates explained 42.91 % of the *nirS*-based denitrifier community changes among all the sampling sites. The distribution of *nirS*-based bacterial community did not show a significant seasonal difference between summer and winter clone libraries (Fig. S2), indicating that the *nirS*-encoding bacterial community compositions were relatively stable in China's coastal wetland sediments. However, a significant latitudinal shift was characterized in the distribution of denitrifier assemblages along the coastal wetlands (Fig. 5).

Quantification of *nirS*-based denitrifiers

The qPCR results indicated a geographically heterogeneous distribution of *nirS*-encoding denitrifier abundance along the coastal wetlands of China (one-way ANOVA, $P < 0.001$) (Fig. 6). The copy numbers of *nirS* gene ranged from 3.1×10^6 to 3.7×10^7 copies g^{-1} dry sediment in summer and 4.3×10^5 to 1.2×10^7 copies g^{-1} dry sediment in winter. The highest copy number was found at the low latitude site P7 in summer, and the lowest copy number was detected at the high latitude site P5 in winter. The average copy number of *nirS*-encoding denitrifying community was slightly higher at the low latitude sites (P7 to P11) than at the high latitude sites (P1 to P6) (Fig. 6). Although no significant seasonal shift was found (one-way ANOVA, $P = 0.079$), the *nirS*-encoding bacterial abundance tended to be higher in summer than in winter (except for sites P6 and P11), with respective average abundance of 9.0×10^6 and 2.9×10^6 copies g^{-1} dry sediment.

Relationships of *nirS*-encoding denitrifier community dynamics with environmental factors

The relationships between the *nirS*-type denitrifier communities with environmental variables were tested by the canonical correspondence analysis (CCA) (Fig. 7). The environmental parameters in the first two CCA dimensions (CCA1 and CCA2) provided 49.9 % of the cumulative variance of the *nirS*-type denitrifying community-environment correlation. The results showed that the *nirS*-based denitrifying community structures in the sediments of China's coastal wetlands were significantly correlated to ammonium ($P = 0.017$, $F = 2.89$, 1000 Monte Carlo permutations) and temperature ($P = 0.048$, $F = 1.63$, 1000 Monte Carlo permutations), which accounted for 37.0 % of the total CCA expositive power. Although the contribution of other investigated environmental variables (including nitrate, nitrite, organic carbon, sulfide, salinity, organic nitrogen, median size, pH, and C:N ratios) were not significant ($P > 0.05$, 1000 Monte Carlo permutations), the union of these environmental factors accounted for additional 45.0 % of the entire CCA expositive power.

The relationships of *nirS*-based bacteria diversity with environmental parameters were also analyzed with SPSS software. The results showed that the diversity of *nirS*-based denitrifying bacteria was positively correlated to organic nitrogen ($R = 0.458$, $P = 0.032$, $N = 22$) and negatively correlated to C:N ratios ($R = -0.540$, $P = 0.009$, $N = 22$) (Table S4). However, no significant correlations were found between *nirS*-based denitrifier diversity and other environmental variables (including temperature, salinity, pH, sediment median size, organic carbon, sulfide, ammonium, nitrite, and nitrate) ($P > 0.05$).

In addition, Pearson correlation analyses indicated that the *nirS*-based denitrifier abundance was only correlated with




Fig. 3 Neighbor-joining phylogenetic tree of *nirS* sequences in China's coastal wetland sediments. This showed affiliations between *nirS* gene fragments derived from the coastal wetlands of China and related matches, with the *Proteobacteria Pseudomonadales* (accession no. D84475) used as the outgroup. Clone names include the sample name

and the number of sequences recovered from each sampling site in summer (red) and winter (blue). Bootstrap values greater than 50 % ($n = 1000$) are shown with solid circle, and those less than 50 % are shown with open circle on the corresponding nodes

temperature ($R = 0.420$, $P = 0.050$), as compared with other environmental variables (Table S4).

Discussion

This study examined the *nirS*-harboring denitrifier community diversity, abundance, and distribution, which provides novel insights into the denitrifier community dynamics and associations with environment factors in the coastal wetlands of China. The diversity of *nirS*-based denitrifiers in this study was consistent with previous reports from other environmental ecosystems (Francis et al. 2013; Li et al. 2013). The majority of detected *nirS* sequences fell into numerous novel phylogenetic lineages and OTUs, most of which might represent coastal wetland-specific *nirS*-encoding denitrifiers. All *nirS* sequences shared considerable phylogenetic similarity with sequences obtained from aquatic systems (marine/estuarine/coastal sediments and hypoxia sea water; Fig. 3), suggesting that all the microorganisms in this study derived from coastal rather than terrestrial environments. The only cultivated sequence is *Proteobacteria Pseudomonadales* (Yan et al. 2008). However, most of the *nirS*-encoding




Fig. 4 The community composition of *nirS*-encoding denitrifiers. S and W in brackets represent summer and winter, respectively




Fig. 5 The UniFrac weighted PCoA analysis of *nirS*-encoding denitrifier communities. S and W represent summer and winter samples, respectively. Red and blue font represent samples from high and low latitude sites, respectively

genotypes observed in the present study were affiliated with uncultured denitrifying strains. Within all the ten clusters, several clusters (II, V, VI, VIII, IX, and X) showed a distinctive latitudinal differentiation along China's coastal wetlands (Table S3), probably suggesting that the diverse denitrifiers have different environmental adaptation strategies. Furthermore, the community structure and distribution of *nirS*-encoding denitrifiers showed obvious latitudinal heterogeneity along the coastal wetlands of China on the basis of statistical analyses (Fig. 5). This result indicates that temperature may be an important environmental parameter shaping the biogeographical distribution and composition of *nirS*-




Fig. 6 The abundance of *nirS*-based denitrifiers in the coastal wetlands of China. Vertical bars show standard error ($n = 3$)




Fig. 7 Canonical correspondence analysis for the correlations of environment factors with the community structure of *nirS*-encoding denitrifiers in both summer (red up-triangle) and winter (blue down-triangle). Temp, OC, C:N, MΦ, NO₃-N, NO₂-N, ON, and NH₄-N represent temperature, organic carbon, C:N ratios, sediment mean size, nitrate, nitrite, organic nitrogen, and ammonium, respectively

based denitrifying communities in the coastal wetlands of China. Furthermore, the latitudinal distribution pattern was also supported by the CCA results (Fig. 7). Previous studies have found that salinity is a dominant environmental variable shaping the biogeographical distribution of denitrifiers (Abell et al. 2010; Francis et al. 2013; Yoshie et al. 2004; Zheng et al. 2015), mainly by comparing different-salinity habitats. However, samples in this study were collected from coastal wetland sediments with relatively consistent salinity. Therefore, we concluded that temperature rather than salinity significantly contributed to the latitudinal distribution of *nirS*-based denitrifiers. Interestingly, a similar distribution pattern was found for the anaerobic ammonium oxidation (anammox) community as well (Hou et al. 2015).

In addition to temperature, other biochemical indices can also influence the distribution and diversity of *nirS*-encoding bacterial communities. It should be noted that a diverse of environmental factors may be significant in shaping the coastal wetland denitrifiers with complex interactions (Dang et al. 2009; Francis et al. 2013; Bulow et al. 2008). In the present study, the ammonium concentration also had significant contribution to *nirS*-based denitrifier community structure, which may be attributed to the increased supply of oxidized nitrogen through nitrification process and thus provides the electron acceptor for denitrification (Avrahami et al. 2002). Similar results have also been observed in the Yangtze estuary and Jiaozhou Bay (Dang et al. 2009; Zheng et al. 2015). Additionally, C:N ratios has been reported to have significant impact on the denitrification activity, nitrite accumulation, and microbial community composition (Her and Huang 1995; Kim et al. 2008; Mosier and Francis 2010). The diversity of

nirS-based denitrifiers was also observed to negatively correlate with C:N ratios in this study. In addition, the *nirS*-based denitrifier diversity was positively affected by organic nitrogen. Therefore, sediment with high nitrogen content may support the coexistence of diverse denitrifying bacteria. Also, the effect of pH on shaping *nirS*-harboring community composition has been documented by Hallin et al. (2009), which is likely due to the narrow pH ranges for optimal growth of microorganisms (Rousk et al. 2010). However, no significant relationship between pH and *nirS*-encoding denitrifier composition was found in the present study. Bulow et al. (2008) have indicated that the greatest control on the diversity of denitrifying communities is environmental stability. Along the coastal wetlands of China, the community of *nirS*-based denitrifying bacteria did not display a statistically significant seasonal shift between summer and winter, indicating that the denitrifiers distribution reflects an adaption to site-specific features.

The abundance of *nirS*-based denitrifiers demonstrated significant spatiotemporal fluctuations in the coastal wetland sediments of China. The copy numbers of *nirS* gene ranged from 4.3×10^5 to 3.7×10^7 copies g^{-1} dry sediment. The values are within the ranges reported in the Colne estuary ($\sim 10^4$ to 10^7 copies per gram of sediment), Chesapeake Bay (2×10^5 to 7×10^7 copies g^{-1} dry sediment), and San Francisco Bay (5.4×10^5 to 5.4×10^7 copies g^{-1} dry sediment) (Bulow et al. 2008; Mosier and Francis 2010; Smith et al. 2007). Similar to the community structure of *nirS*-harboring denitrifiers, *nirS* gene abundance was also observed to be positively correlated with temperature. Braker et al. (2010) reported that temperature exerts a significant effect on the abundance and composition of denitrifying communities in agricultural soil. In general, temperature could either modify the function of existing microorganisms or rebuild microbial communities, and thus changes the fundamental physiologies which drive biogeochemical processes (Schimel and Gullledge 1998). Although it has been reported that multiple environmental factors (e.g., organic matter, pH, nitrate, ammonium, and sediment water) can affect the abundance of *nirS* gene (Dandie et al. 2011; Dong et al. 2009; Kandeler et al. 2006), no significant relationships were found between the *nirS*-encoding denitrifier abundance and other environmental variables detected in this study (including ammonium, C:N ratios, nitrate, nitrite, organic carbon, sulfide, salinity, organic nitrogen, median size, and pH). In addition, denitrifying bacteria abundance tended to be slightly higher in summer than in winter, even though there was no significant seasonal variation in the *nirS* gene abundance. These results further demonstrate the importance of temperature in shaping the dynamics of denitrifiers in the coastal wetlands of China.

Hou et al. (2015) reported that the denitrification rates showed significant latitudinal heterogeneity along the coastal wetlands of China (one-way ANOVA, $p < 0.05$) (Fig. 8), with




Fig. 8 Spatial variation of denitrification rates in sediments. The squares represent means and solid lines represent median values. Boxes enclose the interquartile range, and whiskers show the full range

average rates of $98.82 \mu\text{mol N kg}^{-1} \text{ day}^{-1}$ at the low latitude sites (P7 to P11) and $72.39 \mu\text{mol N kg}^{-1} \text{ day}^{-1}$ at the high latitude sites (P1 to P6). Interestingly, it was observed that the denitrification rates were significantly correlated with the *nirS*-encoding denitrifier abundance ($R = 0.548$, $P < 0.001$) (Fig. S3). The result is consistent with the previous report in the San Francisco Bay estuary (Mosier and Francis 2010). However, the denitrification rates were not significantly related to the denitrifying community composition ($R = -0.126$, $P = 0.57$) (Fig. S4). These relationships imply that the abundance of denitrifiers, more than structure and diversity, predicts the activity of denitrifying community in the coastal wetlands of China.

In conclusion, this study demonstrated the abundance, composition, and distribution of *nirS*-based denitrifying communities along the coastal wetlands of China. To date, the present work represents the most systematic characterization of *nirS*-type denitrifiers based on molecular effort in China's coastal wetland ecosystems. The composition and structure of *nirS*-based denitrifier communities showed distinctive latitudinal heterogeneity along the coastal wetlands of China. The abundance of *nirS* gene varied between 4.3×10^5 and 3.7×10^7 copies g^{-1} dry sediment, with significant spatial heterogeneity. However, there was no significant seasonal shift in the abundance, structure, and distribution of *nirS*-encoding denitrifiers. In this study, temperature was characterized as a key parameter in regulating the latitudinal distribution of denitrifier community abundance, composition, and distribution. This research provides new insights into the dynamics of *nirS*-encoding denitrifiers in China's coastal wetland ecosystems.

Acknowledgments This work was supported by the National Natural Science Foundations of China (Nos. 41322002, 41271114, 41130525, 41071135 and 41501524) and the Program for New Century Excellent Talents in University (NCET), and the China Postdoctoral Science

Foundation (2015 M581567). We thank Wayne S. Gardner and anonymous reviewers for constructive comments on the manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals.

References

- Abell GC, Reville AT, Smith C, Bissett AP, Volkman JK, Robert SS (2010) Archaeal ammonia oxidizers and *nirS*-type denitrifiers dominate sediment nitrifying and denitrifying populations in a subtropical macrotidal estuary. *ISME J* 4:286–300
- Avrahami S, Conrad R, Braker G (2002) Effect of soil ammonium concentration on N₂O release and on the community structure of ammonia oxidizers and denitrifiers. *Appl Environ Microbiol* 68:5685–5692
- Babbin AR, Keil RG, Devol AH, Ward BB (2014) Organic matter stoichiometry, flux, and oxygen control nitrogen loss in the ocean. *Science* 344:406–408
- Boyer EW, Howarth RW, Galloway JN, Dentener FJ, Green PA, Vörösmarty CJ (2006) Riverine nitrogen export from the continents to the coasts. *Glob Biogeochem Cycles* 20:GB1S91
- Braker G, Fesefeldt A, Witzel KP (1998) Development of PCR primer systems for amplification of nitrite reductase genes (*nirK* and *nirS*) to detect denitrifying bacteria in environmental samples. *Appl Environ Microbiol* 64:3769–3775
- Braker G, Ayala-del-Río HL, Devol AH, Fesefeldt A, Tiedje JM (2001) Community structure of denitrifiers, bacteria, and Archaea along redox gradients in Pacific northwest marine sediments by terminal restriction fragment length polymorphism analysis of amplified nitrite reductase (*nirS*) and 16S rRNA genes. *Appl Environ Microbiol* 67:1893–1901
- Braker G, Schwarz J, Conrad R (2010) Influence of temperature on the composition and activity of denitrifying soil communities. *FEMS Microbiol Ecol* 73:134–148
- Bulow SE, Francis CA, Jackson GA, Ward BB (2008) Sediment denitrifier community composition and *nirS* gene expression investigated with functional gene microarrays. *Environ Microbiol* 10:3057–3069
- Canfield DE, Glazer AN, Falkowski PG (2010) The evolution and future of Earth's nitrogen cycle. *Science* 330:192–196
- Cantera JLL, Stein LY (2007) Molecular diversity of nitrite reductase genes (*nirK*) in nitrifying bacteria. *Environ Microbiol* 9:765–776
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Tumbaugh PJ, Walters WA, Widmann J, Yatsunenkov T, Zaneveld J, Knight R (2010) Qiime allows analysis of high-throughput community sequencing data. *Nat Methods* 7:335–336
- Castro-Gonzalez M, Braker G, Farias L, Ulloa O (2005) Communities of *nirS*-type denitrifiers in the water column of the oxygen minimum zone in the eastern South Pacific. *Environ Microbiol* 7:1298–1306
- Coyne MS, Arunakumari A, Averill BA, Tiedje JM (1989) Immunological identification and distribution of dissimilatory heme *cdl* and nonheme copper nitrite reductases in denitrifying bacteria. *Appl Environ Microbiol* 55:2924–2931
- Cui S, Shi Y, Groffman PM, Schlesinger WH, Zhu YG (2013) Centennial-scale analysis of the creation and fate of reactive nitrogen in China (1910–2010). *Proc Natl Acad Sci* 110:2052–2057
- Dalsgaard T, Thamdrup B, Farias L, Revsbech NP (2012) Anammox and denitrification in the oxygen minimum zone of the eastern South Pacific. *Limnol Oceanogr* 57:1331–1346
- Dandie CE, Wertz S, Leclair CL, Goyer C, Burton DL, Patten CL, Zebarth BJ, Trevors JT (2011) Abundance, diversity and functional gene expression of denitrifier communities in adjacent riparian and agricultural zones. *FEMS Microbiol Ecol* 77:69–82
- Dang H, Wang C, Li J, Li T, Tian F, Jin W, Ding Y, Zhang Z (2009) Diversity and distribution of sediment *nirS*-encoding bacterial assemblages in response to environmental gradients in the eutrophied Jiaozhou Bay, China. *Microb Ecol* 58:161–169
- Danovaro R, Gambi C (2002) Biodiversity and trophic structure of nematode assemblages in seagrass systems: evidence for a coupling with changes in food availability. *Mar Biol* 141:667–677
- Dong LF, Smith CJ, Pappaspyrou S, Stott A, Osborn AM, Nedwell DB (2009) Changes in benthic denitrification, nitrate ammonification, and anammox process rates and nitrate and nitrite reductase gene abundances along an estuarine nutrient gradient (the Colne estuary, United Kingdom). *Appl Environ Microbiol* 75:3171–3179
- Earl SR, Valett HM, Webster JR (2006) Nitrogen saturation in stream ecosystems. *Ecology* 87:3140–3151
- Falkowski PG, Fenchel T, DeLong EF (2008) The microbial engines that drive Earth's biogeochemical cycles. *Science* 320:1034–1039
- Francis CA, O'Mullan GD, Cornwell JC, Ward BB (2013) Transitions in *nirS*-type denitrifier diversity, community composition, and biogeochemical activity along the Chesapeake Bay estuary. *Front Microbiol* 4:237
- Galloway JN, Cowling EB (2002) Reactive nitrogen and the world: 200 years of change. *Ambio* 31:64–71
- Galloway JN, Dentener FJ, Capone DG, Boyer EW, Howarth RW, Seitzinger SP, Asner GP, Cleveland CC, Green PA, Holland EA, Karl DM, Michaels AF, Porter JH, Townsend AR, Osmarty CJV (2004) Nitrogen cycles: past, present, and future. *Biogeochemistry* 70:153–226
- Galloway JN, Townsend AR, Erisman JW, Bekunda M, Cai Z, Freney JR, Martinelli LA, Seitzinger SP, Sutton MA (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320:889–892
- Gruber N, Galloway JN (2008) An earth-system perspective of the global nitrogen cycle. *Nature* 451:293–296
- Hallin S, Jones CM, Schloter M, Philippot L (2009) Relationship between N-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment. *ISME J* 3:597–605
- Hannig M, Braker G, Dippner J, Jurgens K (2006) Linking denitrifier community structure and prevalent biogeochemical parameters in the pelagial of the Central Baltic proper (Baltic Sea). *FEMS Microbiol Ecol* 57:260–271
- He Q, Bertness MD, Bruno JF, Li B, Chen G, Coverdale TC, Altieri AH, Bai J, Sun T, Pennings SC, Liu J, Ehrlich PR, Cui B (2014) Economic development and coastal ecosystem change in China. *Sci Rep* 4:5995
- Her J, Huang J (1995) Influences of carbon source and C/N ratio on nitrate/nitrite denitrification and carbon breakthrough. *Bioresour Technol* 54:45–51
- Hou L, Zheng Y, Liu M, Li X, Lin X, Yin G, Gao J, Deng F, Chen F, Jiang X (2015) Anaerobic ammonium oxidation and its contribution to nitrogen removal in China's coastal wetlands. *Sci Rep* 5:15621
- Howarth RW (2008) Coastal nitrogen pollution: a review of sources and trends globally and regionally. *Harmful Algae* 8:14–20
- Kandeler E, Deiglmayr K, Tschirko D, Bru D, Philippot L (2006) Abundance of *narG*, *nirS*, *nirK*, and *nosZ* genes of denitrifying bacteria during primary successions of a glacier foreland. *Appl Environ Microbiol* 72:5957–5962

- Kim M, Jeong SY, Yoon SJ, Cho SJ, Kim YH, Kim MJ, Ryu EY, Lee SJ (2008) Aerobic denitrification of *Pseudomonas putida* AD-21 at different C/N ratios. *J Biosci Bioeng* 106:498–502
- Kumar S, Tamura K, Nei M (2004) MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* 5:150–163
- Li M, Hong YG, Cao HL, Gu JD (2013) Community structures and distribution of anaerobic ammonium oxidizing and *nirS*-encoding nitrite-reducing bacteria in surface sediments of the South China Sea. *Microb Ecol* 66:281–296
- Lin T, Xue XZ, Lu CY (2007) Analysis of coastal wetland changes using the "DPSIR" model: a case study in Xiamen, China. *Coast Manag* 35:289–303
- Mohamed NM, Saito K, Tal Y, Hill RT (2010) Diversity of aerobic and anaerobic ammonia-oxidizing bacteria in marine sponges. *ISME J* 4:38–48
- Mosier AC, Francis CA (2010) Denitrifier abundance and activity across the San Francisco Bay estuary. *Environ Microbiol Rep* 2:667–676
- Rousk J, Baath E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer N (2010) Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* 4:1340–1351
- Schimel JP, Gullede J (1998) Microbial community structure and global trace gases. *Glob Chang Biol* 4:745–758
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75:7537–7541
- Seitzinger S (2008) Nitrogen cycle: out of reach. *Nature* 452:162–163
- Shi L, Liu F, Zhang Z, Zhao X, Liu B, Xu J, Wen Q, Yi L, Hu S (2015) Spatial differences of coastal urban expansion in China from 1970s to 2013. *Chin Geogr Sci* 25:389–403
- Smith CJ, Nedwell DB, Dong LF, Osborn AM (2007) Diversity and abundance of nitrate reductase genes (*narG* and *napA*), nitrite reductase genes (*nirS* and *nrfA*), and their transcripts in estuarine sediments. *Appl Environ Microbiol* 73:3612–3622
- Stauffert M, Cravo-Laureau C, Duran R (2014) Structure of hydrocarbonoclastic nitrate-reducing bacterial communities in bioturbated coastal marine sediments. *FEMS Microbiol Ecol* 89:580–593
- Sun Z, Sun W, Tong C, Zeng C, Yu X, Mou X (2015) China's coastal wetlands: conservation history, implementation efforts, existing issues and strategies for future improvement. *Environ Int* 79:25–41
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599
- ter Braak CJF (1988) CANOCO: an extension of DECORANA to analyze species-environment relationships. *Vegetatio* 75:159–160
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882
- Throckmold IN, Enwall K, Jarvis A, Hallin S (2004) Reassessing PCR primers targeting *nirS*, *nirK* and *nosZ* genes for community surveys of denitrifying bacteria with DGGE. *FEMS Microbiol Ecol* 49:401–417
- van Breemen N, Boyer EW, Goodale CL, Jaworski NA, Paustian K, Seitzinger SP, Lajtha K, Mayer B, van Dam D, Howarth RW, Nadelhoffer KJ, Eve M, Billen G (2002) Where did all the nitrogen go? Fate of nitrogen inputs to large watersheds in the northeastern USA. *Biogeochemistry* 57:267–293
- Vitousek PM, Aber JD, Howarth RW (1997) Human alteration of the global nitrogen cycle: sources and consequences. *Ecol Appl* 7:737–750
- Wang Y (1992) Coastal management in China. In: Fabbri P (ed) *Ocean coastal management in global change*. Elsevier, London, pp. 460–469
- Wang XM, Wang JL (2009) Removal of nitrate from groundwater by heterotrophic denitrification using the solid carbon source. *Sci China, Ser B: Chem* 52:236–240
- Wetland China (2014) The report on the second national wetland resources survey (2009–2013). <http://www.shidi.org/zt/2014xwfbh/>
- Xie Z, Xu X, Yan L (2010) Analyzing qualitative and quantitative changes in coastal wetland associated to the effects of natural and anthropogenic factors in a part of Tianjin, China. *Estuar Coast Shelf Sci* 86:379–386
- Yan Y, Yang J, Dou Y, Chen M, Ping S, Peng J, Lu W, Zhang W, Yao Z, Li H, Liu W, He S, Geng L, Zhang X, Yang F, Yu H, Zhan Y, Li D, Lin Z, Wang Y, Elmerich C, Lin M, Jin Q (2008) Nitrogen fixation island and rhizosphere competence traits in the genome of root-associated *Pseudomonas stutzeri* A1501. *Proc Natl Acad Sci U S A* 105:7564–7569
- Yoshie S, Noda N, Tsuneda S, Hirata A, Inamori Y (2004) Salinity decreases nitrite reductase gene diversity in denitrifying bacteria of wastewater treatment systems. *Appl Environ Microbiol* 70:3152–3157
- Zhang Y, Xie X, Jiao N, Hsiao SY, Kao SJ (2014) Diversity and distribution of *amoA*-type nitrifying and *nirS*-type denitrifying microbial communities in the Yangtze River estuary. *Biogeosciences* 11:2131–2145
- Zhao X, Zhou Y, Min J, Wang S, Shi W, Xing G (2012) Nitrogen runoff dominates water nitrogen pollution from rice-wheat rotation in the Taihu Lake region of China. *Agric Ecosyst Environ* 156:1–11
- Zheng Y, Hou L, Liu M, Gao J, Yin G, Li X, Deng F, Lin X, Jiang X, Chen F, Zong H, Zhou J (2015) Diversity, abundance, and distribution of *nirS*-harboring denitrifiers in intertidal sediments of the Yangtze estuary. *Microb Ecol* 70:30–40
- Zumft WG (1997) Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 61:533–616